New species and combinations of *Inocybe* with lilac or violet colours in Europe

MUÑOZ G.¹, PANCORBO F.², TURÉGANO Y.³, & ESTEVE-RAVENTÓS F.⁴

¹ Calle Tudela, 20, 50650 Gallur, Zaragoza, Spain (Grupo Cultural Micológico Verpa). E-mail: guillermomunoz1981@gmail.com

² Sociedad Micológica de Madrid, Real Jardín Botánico. C/ Claudio Moyano 1, 28014 Madrid, Spain. E-mail: fpancorbo@gmail.com

³ Departamento de Ciencias de la Vida (Área de Botánica). Universidad de Alcalá, 28805 Alcalá de Henares, Madrid, Spain. E-mail: yolanda.turegano@edu.uah.es

⁴ Departamento de Ciencias de la Vida (Área de Botánica). Universidad de Alcalá, 28805 Alcalá de Henares, Madrid, Spain. E-mail: fernando.esteve@uah.es

Resumen

Se revisa el grupo de especies con apariencia y hábito de *I. pusio*, todas con un estípote más o menos lilacino a violáceo, sobre todo en ejemplares jóvenes, mediante el estudio de caracteres morfológicos y análisis de la región ITS. La distribución de los caulocistidios, la presencia de vellipellis y la forma esporal son los caracteres más importantes que permiten la identificación de especies los cuales se pudieron correlacionar con la señal filogenética de la región ITS. Se obtuvieron clados en el nivel de especie bien apoyados en los análisis filogenéticos, salvo para *I. paleoveneta*. Como resultado, se describen tres especies nuevas: *Inocybe caesaraugustae*, *I. ianthinopes* e *I. velatipusio*, y se proponen dos combinaciones nuevas: *I. floccipes* e *I. paleoveneta*, poniéndose de manifiesto la importancia del estudio de la distribución de los caulocistidios en este grupo. Una característica común a varias de las especies tratadas (*I. caesaraugustae*, *I. ianthinopes*, *I. pusio*, *I. paleoveneta*) es su fructificación habitual en áreas con cierta influencia humana en ambientes ruderales y suelos nitrificados, como parques y jardines. Se incluyen descripciones e ilustraciones de todas las especies presentadas.

Palabras clave: *Inocybe* sect. *Lilacinae*, *Inocybe pusio*, taxonomía, morfología, distribución geográfica, filogenia, Europa.

Resumo

É feita uma revisão do grupo de espécies com aparência de *Inocybe pusio*, todos com um estipe mais ou menos lilacino a violáceo, sobretudo em exemplares jovens, mediante o estudo de caracteres morfológicos e análise da região ITS. A distribuição dos caulocistídios, a presença de vellipellis e a forma dos esporos são os caracteres mais importantes que permitem a identificação de espécies e podem ser correlacionados com o análise filogenética da região ITS. Foram obtidos clados bem suportados nas análises filogenéticas, excepto para *I. paleoveneta*. Como resultado, são descritas três espécies novas: *Inocybe caesaraugustae*, *I. ianthinopes* e *I. velatipusio*, e propostas duas combinações novas: *I. floccipes* e *I. paleoveneta*, sendo posta em evidência a importância do estudo da distribuição dos caulocistídios neste grupo. Uma característica comum a algumas das espécies tratadas (*I. caesaraugustae*, *I. ianthinopes*, *I. pusio*, *I. paleoveneta*) é a frutificação habitual em áreas com alguma influência humana em ambiente ruderal e solos nitrificados, como parques e jardins. Estão incluídas descrições e ilustrações de todas as espécies apresentadas.

Palavras chave: *Inocybe* sect. *Lilacinae*, *Inocybe pusio*, taxonomia, morfologia, distribuição geográfica, filogenia, Europa.

Summary

The group of species with the appearance and habit of *I. pusio*, all with a more or less lilac to purplish stipe, especially in young specimens, is reviewed based on the study of morphological characters and analyses of the ITS region. The distribution of the caulocystidia, the presence of vellipellis, and the spore shape are the most important characters allowing for species identification, and were correlated with the phylogenetic signal of the ITS region. Well-supported species-level clades were obtained in the phylogenetic analyses, except for *I. paleoveneta*. As a result, three new to science species are described: *Inocybe caesaraugustae*, *I. ianthinopes* and *I. velatipusio*, and two new combinations are proposed: *I. floccipes* and *I. paleoveneta*, highlighting the importance of studying the distribution of caulocystidia in this group. A common feature of several of the species discussed (*I. caesaraugustae*, *I. ianthinopes*, *I. pusio*, *I. paleoveneta*) is their occurrence in places with some human influence, often in ruderal areas and nitrified soils, such as parks and gardens. Descriptions and illustrations of all the species presented are included.

Key-words: *Inocybe* sect. *Lilacinae*, *Inocybe pusio*, taxonomy, morphology, geographic distribution, phylogeny, Europe.

Introduction

In the genus *Inocybe* (Fr.) Fr. it is relatively common to find in Europe a good number of species showing lilac, violet or wine-purple shades in some part of the basidiomata, especially on the stipe, but also more infrequently on the pileus and lamellae; this chromatic particularity, so attractive for the genus, was the reason why HEIM (1931) grouped all of them in *I. sect. Lilacinae* R. Heim. According to HEIM (1931: 250), the species comprising *I. sect. Lilacinae* are characterised not only by the particular colours, but also by the presence of a cortina and smooth spores. BON (1997) selected *I. pusio* P. Karst. as the type of this section.

The taxonomic recognition and practical use of Heim's *I. sect. Lilacinae* was commonly accepted in subsequent years in Europe (KÜHNER & ROMAGNESI 1953; STANGL & VESELSKÝ 1982; MOËNNE-LOCCOZ & *al.* 1988; BON 1997), although the species traditionally included in *I. sect. Lilacinae* differ in appearance, colour of the basidiomata, surface of the pileus, reaction of the cystidial walls in ammoniacal media and, above all, the distribution of the caulocystidia. These facts were already commented by BON (1997: 8), although he and CARTERET & REMAUX (2011) maintained *I. sect. Lilacinae* as practical grouping, in spite of its marked polyphyletism.

In modern classifications and keys of the genus *Inocybe* *s.l.*, *I. sect. Lilacinae* has been progressively abandoned and well-known species [e.g. *I. pusio*, *I. cincinnata* (Fr.) Quél. or *I. griseolilacina* J.E. Lange] have been included in more natural phyletic groups, and integrated in different groups or sections such as *I. sect. Fibrillosae* R. Heim, *I. sect. Tardae* Bon, *I. sect. Cortinatae* Kühner & Boursier, etc. (JACOBSON 2008; JACOBSON & LARSSON 2012). It can be stated that in practically every section or taxonomic group of the genus *Inocybe* one can find species with lilac or purplish hues; these particular colours can also be present in taxa both with smooth or nodulose spores (e.g. *I. lacera* (Fr.) P. Kumm. *s.l.*, *I. curvipes* P. Karst. *s.l.*, *I. putilla* Bres., or *I. gonio-pusio* Stangl).

Recently, according to an extensive phylogenetic study and the current recognition of seven genera within the family *Inocybaceae* Jülich (MATHENY & *al.* 2020), the genus *Inosperma* (Kühner) Matheny & Esteve-Rav. also includes species with violet to lilac colours, e.g. *I. insignissimum* (Romagn.) Matheny & Esteve-Rav., *I. lilofastigiatum* (Stangl & J. Veselský) Matheny & Esteve-Rav., *Inocybe quietiodor* f. *lilacino-pes* Carteret & Reumaux, etc.

As a result of our studies, we assembled a good set of material of species similar to *I. pusio*, among which several collections did not conform to *I. pusio*

s. str. Thus, the main objectives of this study were to i) assess species limits among species resembling *I. pusio* based on molecular, morphological and ecological characters and ii) to propose an updated nomenclature for the group.

Materials and methods

Morphological and material study

The studied collections have been photographed macroscopically *in situ* with a Nikon D50 and an Olympus OMD E-M5 MarkII digital cameras, using tripod and natural light. The macroscopic descriptions are based on fresh material, which was subsequently dehydrated for herbarium conservation. Water and 10-30% ammonia $\text{NH}_3(\text{aq})$ were used for microscopic preparations. For the microscopic observations and their corresponding descriptions, a Motic BA300 optical microscope with a Moticam microphotographic camera or Nikon 55i microscope with a Nikon D90 camera, connected to a personal computer, was used. Subsequently, the images taken were processed with a computer program for images (Adobe Photoshop or Lightroom). The measurements of spores and cystidia were done using Piximètre 5.10 R 1541, the "classique" format has been chosen to present the results (<http://ach.log.free.fr/Piximetre/Piximetre5.htm>). Cystidia were measured without crystals and basidia without sterigmata.

The descriptions, both macroscopic and microscopic, were always those observed by the authors in their collections. The material has been deposited in the herbarium of the University of Alcalá (AH) and in the private herbarium of some of the authors (Guillermo Muñoz, indicated as GM and Fermín Pancorbo indicated as FP).

Colour codes are taken from MUNSELL (1994), terminology follows VELLINGA (1988) and KUYPER (1986).

Abbreviations and symbols

- Cl_3Hg : trichloromercury(1-)
- L_m : average length
- L: number of lamellae reaching from the margin of the pileus to the stipe
- l: number of lamellae not reaching the stipe
- n: number of spores or structures measured for the calculation of statistics
- N: number of collections
- $\text{NH}_3(\text{aq})$: ammonia aqueous solution
- Q: ratio of spore length to spore width
- Q_m : average ratio L_m / W_m
- W_m : average width

Molecular study

The aim of the molecular studies performed is to support the cited material through a molecular interpretation. DNA extractions were performed from dried material using a NZY Plant/Fungi gDNA Isolation Kit (NZYTech, Lisboa, Portugal). The amplified region was the ITS rDNA, which was amplified using primers ITS5 and ITS4 (WHITE & *al.* 1990). PCR conditions were as follows: 5 min at 95 °C; 38 cycles of 45 s at 95 °C, 45 s at 52 °C, 1 min at 72 °C; 10 min at 72 °C. Fragments were visualised by electrophoresis on 1% agarose gels, stained with SYBR Safe DNA Gel Stain (Invitrogen by Thermo Fisher Scientific) and visualised with a blue light transilluminator. The amplified samples were purified by precipitation with 4M ammonium acetate, isopropanol and 70% ethanol. Amplicons were sequenced at the Molecular Biology Unit (Faculty of Environmental Sciences, University of Alcalá de Henares) using primers ITS2, ITS3, ITS4 and ITS5 (WHITE & *al.* 1990). The consensus ITS sequence of the four primers was obtained using Geneious 5.1.7 (<https://www.geneious.com>). To preliminarily verify the identity of the sequences, BLAST searches were performed (ALTSCHUL & *al.* 1990).

A total of 15 sequences of the ITS region were generated in this work (Table 1). Alignments were assembled by downloading the closest records in BLAST searches and sequences from both *Nothocybe distincta* (K.P.D. Latha & Manim.) Matheny & K.P.D. Latha and *Inocybe lacera* (Fr.) P. Kumm. were used as outgroups to root the tree. Sequences were automatically aligned in Aliview (LARSSON 2014) and then adjusted manually. All characters of the ITS region could be aligned unambiguously. As the phylogenetic inferences did not differ when the ITS1, 5.8S and ITS2 regions were partitioned, the analyses were run without any partition not to over-parameterize them. Alignments were analyzed using maximum likelihood (ML) and Bayesian inference (BI) methods. ML analyses were performed in IQ-TREE web server (TRIFINOPOULOS & *al.* 2016), leaving all other options as default. To determine node support, 1000 ultrafast bootstrap replicates were performed. The substitution model was calculated by ModelFinder implemented in IQ-TREE web server. The best fitting model for ML analysis was HKY+F+G4. MB analysis was performed in MrBayes 3.2.7 (RONQUIST & *al.* 2012). The substitution models were sampled within GTR space (RONQUIST & *al.* 2012), using two analyses of four MCMC chains over 5 million generations, starting with a random tree and sampling one tree every 1000 generations. Chains were assumed to converge properly when the standard deviation values of cluster frequencies fell below 0.01 (1016000 generations).

A burn-in of 25% of the trees was selected and discarded from each analysis. Branch support (posterior probability values, BPP) was calculated using the consensus tree following the “50% majority rule” of the remaining trees. Nodes with ML bootstrap value >95% and BPP ≥0.95 were considered phylogenetically supported.

Results

The most likely tree of the Maximum Likelihood analysis, presented in Fig. 1, and the 50% majority rule consensus tree from the Bayesian analysis showed a similar topology, but the latter analysis yielded more supported nodes. Nearly all the species treated in this article nest in well-supported clades in both analyses, such as *Inocybe cf. pusio* (ML-BS 100%, BPP 1), *I. ianthinopes* (ML-BS 100%, BPP 1), *I. floccipes* (ML-BS 96%, BPP 1), and *I. velatipusio* (ML-BS 100%, BPP 1). Exceptions were *I. caesaraugustae*, not supported in the Maximum Likelihood analysis (ML-BS 94%, BPP 1), and *I. paleoveneta*, not supported in any analyses (ML-BS 76%, BPP 0.66). Nevertheless, all sequences of *I. paleoveneta* included in our analyses are identical and form a distinct clade divergent from that of *I. caesaraugustae*.

The backbone of the tree is fully supported in the Bayesian analysis and the *I. paleoveneta*-*I. caesaraugustae* clade is furthermore supported in the Maximum Likelihood analysis (ML-BS 100%, BPP 1).

Taxonomy

Inocybe caesaraugustae G. Muñoz, Esteve-Rav. & Pancorbo, *sp. nov.*

Mycobank: MB 841753

Holotypus: SPAIN. **Zaragoza**: Zaragoza, Parque Grande José Antonio Labordeta, Jardín Botánico, 41.633972 -0.895444 (±10 m), 243 m, on muddy soil, under *Tilia tomentosa*, *Prunus laurocerasus*, *Arbutus unedo* and *Berberis vulgaris*, 25-X-2012, leg. G. Muñoz, AH 56200 (ITS GenBank OL352083). *Isotypus* in GM2677.

Figs. 2-3

Etymology: *caesaraugustae* (Latin), in reference to the Roman name of the city of Zaragoza (*Caesaraugusta*), where the material was collected.

Diagnosis: Similar morphologically to *I. paleoveneta* but differing by its ellipsoidal to subamygdaliform spores with rounded rather than pointed-ovoid apex and slightly wider cystidial walls (1.5-2.5 µm

Table 1. ITS sequences of the taxa employed in the study, indicating accepted name and original (in brackets) identifications, voucher numbers, geographical origin, possible hosts and GenBank accession numbers. Bold entries refer to newly generated sequences. * means ITS/LSU sequences

Identification (<i>I.</i> = <i>Inocybe</i>)	Voucher specimen	Geographical origin	Possible host	ITS GenBank accession number
<i>I. caesaraugustae</i>	AH 40372	Spain	<i>Tilia tomentosa</i> , <i>Prunus laurocerasus</i> , <i>Arbutus unedo</i> , <i>Berberis vulgaris</i>	OL352085
<i>I. caesaraugustae</i>	AH 56200 holotype	Spain	<i>Tilia tomentosa</i> , <i>Prunus laurocerasus</i> , <i>Arbutus unedo</i> , <i>Berberis vulgaris</i>	OL352083
<i>I. caesaraugustae</i>	AH 40371	Spain	<i>Tilia tomentosa</i> , <i>Prunus laurocerasus</i> , <i>Arbutus unedo</i> , <i>Berberis vulgaris</i>	OL352084
<i>I. caesaraugustae</i> (<i>I.</i> aff. <i>grammopodia</i>)	M72	Hungary	<i>Tilia tomentosa</i>	MW354973
<i>I. derbschii</i>	PRM716232 paratype of <i>I. pseudodestructa</i>	Germany	<i>Pinus</i>	MG012469
<i>I. derbschii</i>	DB30-5-13-2	Germany	<i>Picea abies</i>	MH366593
<i>I. floccipes</i> (<i>I. pusio</i> var. <i>floccipes</i>)	AH 24988	Spain	<i>Fagus sylvatica</i>	OL352058*
<i>I. floccipes</i> (<i>I. pusio</i> var. <i>floccipes</i>)	AH 29885	Portugal	<i>Quercus faginea</i>	OL352059*
<i>I. floccipes</i> (<i>I. pusio</i> var. <i>floccipes</i>)	AH 40051	Italy	<i>Quercus</i>	OL352061*
<i>I. floccipes</i> (<i>I. pusio</i> var. <i>floccipes</i>)	AH 46767	Spain	<i>Quercus ilex</i> subsp. <i>ilex</i>	OL352060*
<i>I. floccipes</i> (<i>I. pusio</i> var. <i>floccipes</i>)	AH 30626 holotype	Spain	<i>Fagus sylvatica</i>	OK414030
<i>I. ianthinopes</i>	AH 40370 holotype	Spain	<i>Tilia tomentosa</i> , <i>Prunus laurocerasus</i> , <i>Arbutus unedo</i> , <i>Berberis vulgaris</i>	OL352086
<i>I. ianthinopes</i> (<i>I.</i> cf. <i>pusio</i>)	AH 48228	Spain	<i>Corylus avellana</i>	OL352087
<i>I. ianthinopes</i> (<i>I. griseolilacina</i>)	Bizio 16919	Italy	<i>Fraxinus</i> , <i>Populus</i>	JF908164
<i>I. ianthinopes</i> (<i>I. pusio</i>)	DB16-8-14-24	Germany	Unknown	MH366588
<i>I. ianthinopes</i> (<i>I. pusio</i>)	DB14-7-12-4	Germany	Unknown	MH366589
<i>I. paleoveneta</i>	AH 51448	Spain	<i>Carpinus betulus</i>	OL583808*

Identification (<i>I.</i> = <i>Inocybe</i>)	Voucher specimen	Geographical origin	Possible host	ITS GenBank accession number
<i>I. paleoveneta</i> (<i>I.</i> cf. <i>grammopodia</i>)	KR-M-0044789	Germany	<i>Quercus robur</i>	MT005886
<i>I. paleoveneta</i> (<i>I. grammopodia</i> var. <i>paleoveneta</i>)	MCVE:29415 holotype	Italy	<i>Betula pendula</i> , <i>Carpinus betulus</i>	NR_164491
<i>I. paleoveneta</i> (<i>I. grammopodia</i>)	IMG 1632	USA	Unknown	KY680785
<i>I. paleoveneta</i> (<i>I. grammopodia</i>)	KR-M-0044812	Germany	<i>Quercus cerris</i>	MT005892
<i>I. paleoveneta</i> (<i>I. grammopodia</i>)	KR-M-0044138	Germany	<i>Tilia cordata</i>	MH366590
<i>I. paleoveneta</i> (<i>I. grammopodia</i>)	DB14-10-15-2	Germany	Unknown	MH366591
<i>I.</i> cf. <i>pusio</i> (<i>I. pusio</i>)	(TUR-A) JV27197F	Estonia	<i>Tilia</i> , <i>Quercus robur</i> , <i>Betula</i>	KJ432288
<i>I.</i> cf. <i>pusio</i> (<i>I. pusio</i>)	(TUR-A) JV17838	Unknown	Unknown	KJ432287
<i>I.</i> cf. <i>pusio</i> (<i>I. pusio</i>)	(TUR-A) 78010	Sweden	<i>Fraxinus</i> , <i>Corylus</i>	KJ432286
<i>I.</i> cf. <i>pusio</i> (<i>I. pusio</i>)	AH 40053	Italy		OL352057*
<i>I.</i> cf. <i>pusio</i> (<i>I. amethystina</i>)	21605	Italy	Unknown	JF908241
<i>I. velatipusio</i>	AH 47725 holotype	France	<i>Corylus avellana</i> , <i>Fagus sylvatica</i> , <i>Abies alba</i>	OL352088
<i>I. velatipusio</i>	AH 34397	Spain	<i>Quercus faginea</i>	OL352089
<i>I. velatipusio</i>	AH 45163	Spain	<i>Quercus faginea</i> , <i>Cistus monspeliensis</i>	OL352090
<i>I. velatipusio</i> (Uncultured fungus)	clone 23	France	<i>Quercus pubescens</i>	FN397124
<i>I. velatipusio</i> (Uncultured <i>Inocybe</i>)	clone 39	Iran	<i>Fagus orientalis</i> , <i>Carpinus betulus</i> , <i>Quercus castaneifolia</i> , <i>Q. macranthera</i>	FR852257
<i>I. lacera</i>	EL-2104	Sweden		AM882823
<i>Nothocybe distincta</i>	CAL 1310	India		NR_173156

Fig. 1. Most probable tree inferred by maximum likelihood (ML) analysis of the ITS region of the rDNA in species related to *Inocybe pusio*. Bootstrap-ML values / posterior probability from Bayesian analysis around the branches are shown. Thick branches indicate nodes with phylogenetic support in both analysis (bootstrap values $\geq 95\%$ and posterior probability ≥ 0.95). Sequences of *Inocybe lacera* and *Nothocybe distincta* to root the tree. The country of origin of each collection is abbreviated by ISO codes, with specimens described in this article marked in bold.

thick). *Inocybe pusio* shows a different distribution of the caulocystidia and *I. ianthinopes* a different pileus appearance, woolly-fibrillose in the latter. It differs phylogenetically from the closest species, *I. paleoveneta*, by 23 bp in the ITS region (Blast 96.4 % of similarity).

Basidiomata agaricoid and stipitate. *Pileus* 10-35(-40) mm diam, at first hemisphaerical to conico-convex, then plano-convex, obtusely umbonate in adult specimens, not hygrophanous; margin involute when young, regular to wavy in old specimens, surface covered by a thin white to greyish-white velipellis in young specimens, in some collections more or

Fig. 2. *Inocybe caesaraugustae*. A: Collection AH 56200. Holotype. B: Collection AH 40371. C: Collection AH 40372. Scale bar: 20 mm.

less persistent, in others fleeting; colour uniformly chesnut brown in young basidiomata, becoming leather brown with age, at least towards the margin which can be paler with development (Mu 7.5YR 5/4-6, at the centre 7.5YR 4/4); surface unevenly fibrillose at first, then floccose to furfuraceous at the center, often cracked concentrically into appressed scales (“tabby”, reminding *I. subtigrina* Kühner), radially fibrillose at the extreme margin, dry. *Lamellae* moderately crowded ($L = 40-52$; $l = 0-1$), adnexed to emarginate, ventricose, initially lilac grey with

glaucous reflections, becoming lilac to pale brown or greyish brown, finally tobacco-brown, edge paler to concolourous, finely fimbriate. *Stipe* 20-50 × 2-5 mm, straight to frequently curved or sinuose, cylindrical or hardly enlarged towards the base; colour intensely and uniformly violaceous to blue-violaceous at first, becoming lilac ochraceous at the the base with age; surface often striated longitudinally, smooth or covered with whitish fibrils depending on development, fibrillose-floccose at the extreme apex near the pileus. *Cortina* present. *Context* fibrose, whitish and scanty

Fig. 3. *Inocybe caesaraugustae* AH 56200. Holotype. A: Basidiospores. B: Cheilocystidia. C: Pleurocystidia. D: Caulocystidia and caulocystidioid hairs. Scale bar: A: 10 μm , B-D: 50 μm . Mounting media: A-D: $\text{NH}_3(\text{aq})$.

at the pileus, lilac or pale violaceous at the stipe. *Smell* strongly spermatic, taste not significant.

Basidiospores smooth, yellowish ochraceous, ellipsoid to subamygdalaform with rounded apex, sometimes conical, slightly thickened walls ($\approx 0.5\mu\text{m}$) of $(7.7\text{--}8.3\text{--}10.4\text{--}11.6) \times (4.8\text{--}5.1\text{--}6.2\text{--}6.7)$, $Q = (1.4\text{--}1.5\text{--}1.8\text{--}2.1)$; ($L_m=9.3\mu\text{m}$, $W_m=5.7\mu\text{m}$, $Q_m=1.6$; $n=172$, $N=3$). **Basidia** hyaline, claviform, 4-spored, very occasionally 2-spored, $(25.3\text{--})26.5\text{--}34.3\text{--}(35.7) \times (6.8\text{--})7.9\text{--}10.2\text{--}(12.2)\mu\text{m}$, ($L_m=30.8\mu\text{m}$, $W_m=9.2\mu\text{m}$, $n=40$). **Cheilocystidia** $(37\text{--})45\text{--}68\text{--}(93) \times (9.6\text{--})13\text{--}19\text{--}(56)\mu\text{m}$, ($L_m=56\mu\text{m}$, $W_m=17\mu\text{m}$), very abundant, metuloid, morphologically variable, mostly lageniform and ventricose, to subcylindrical or subfusiform, or sublageniform with long and often wavy neck, thick walled $(1\text{--})1.3\text{--}2.6\text{--}(3.5)\mu\text{m}$, $W_m = 1.8\mu\text{m}$, slightly yellowish in $\text{NH}_3(\text{aq})$; **paracystidia** intermixed with cheilocystidia, sometimes very abundant, especially near the pileus margin, subhyaline or filled with yellowish content, claviform to pyriform, $15\text{--}28 \times 6\text{--}17\mu\text{m}$ ($L_m=22\mu\text{m}$, $W_m=11\mu\text{m}$), often articulated at the base, some thick-walled ($\approx 1.5\mu\text{m}$). **Pleurocystidia** $(43\text{--})46\text{--}64\text{--}(105) \times (8.8\text{--})11\text{--}18\text{--}(59)\mu\text{m}$; ($L_m=56\mu\text{m}$, $W_m=15\mu\text{m}$, $n=74$, $N=3$) similar to cheilocystidia, also morphologically variable, with

predominance of fusiform to lageniform, occasionally septe and filled with yellow ochraceous content. **Stipitipellis** consisting of a cutis of parallel hyphae of $(3.7\text{--})4.7\text{--}8.8\text{--}(13)\mu\text{m}$ ($W_m=6.9$); metuloid **caulocystidia** present only in the apical part of the stipe (in upper 1/10 of stipe) of $(35)45\text{--}68\text{--}(79) \times (8.3\text{--})10\text{--}17\text{--}(21)\mu\text{m}$ ($L_m=56\mu\text{m}$, $W_m=13\mu\text{m}$), apparently absent in some collections (AH 40372), mixed with numerous caulocystidioid hairs, $7\text{--}10\mu\text{m}$ wide. **Clamps** present.

Habitat & Distribution: It occurs in more or less numerous groups on moist soil under various broad-leaved trees. The material studied was collected in ruderalised and nitrified areas, such as parks and gardens, on calcareous soils. A common feature between the samples from Zaragoza and that of the reference GB MW354973 (as *I. aff. grammopodia*, voucher M72) from Hungary, is the presence of silver lime trees (*Tilia tomentosa*) in their vicinity (CZISMAR & al. 2021), although this host and kind of habitat may also be usual in *I. paleoveneta* (see comments on this species) and *I. pusio* (LUDWIG 2017).

Additional material studied: SPAIN. **Zaragoza:** Zaragoza, Parque Grande José Antonio Laborde-ta, Jardín Botánico, $41.633972\text{--}0.895444 (\pm 10\text{ m})$,

243 m, on muddy soil, under *Tilia tomentosa*, *Prunus laurocerasus*, *Arbutus unedo* and *Berberis vulgaris*, 28-X-2009, leg. G. Muñoz, AH 40371, duplicate in GM1647 (ITS GenBank OL352084); *ibidem*, 10-XI-2009, leg. G. Muñoz, AH 40372, duplicate in GM1703 (ITS GenBank OL352085).

Comments: The similarity between *I. caesaraugustae* and *I. grammopodia* var. *paleoveneta* Bizio & A. Castellan is evident, both in morphological and ecological characters. However, our analyses (Fig. 1) show that specimens of *I. caesaraugustae* nest in a supported clade in the Bayesian analysis, whereas the specimens of *I. grammopodia* var. *paleoveneta* appear outside that clade (Fig. 1). According to our analyses, both species are, however, close as they fall within a more comprehensive fully supported clade. The collection from Hungary under *Tilia* (as *I. aff. grammopodia*, CZISMAR & al. 2021) falls into the same clade as *I. caesaraugustae*, whereas those obtained from GenBank, identified as *I. grammopodia* Malençon from Germany, correspond to *I. paleoveneta*.

Inocybe paleoveneta, as originally described by BIZIO & CASTELLAN (2018, sub *I. grammopodia* var. *paleoveneta*), differs macro- and microscopically from *I. caesaraugustae*. *Inocybe paleoveneta* was described as having a rimose pileus at the margin with development, with an abundant veil when young, the colour of the stipe is never indicated as violet (it is described with a lilac to lilac-pink hue), and there is no mention that the lamellae show a lilac colour. Furthermore, the spores of *I. paleoveneta* show a slight suprahilar depression and their apex is conical to ogival, sometimes even papillate, and besides, the cystidia have (not always) a distinct neck and cystidial walls are reported as very thin (0.5–1 µm). All these characters differ from those of *I. caesaraugustae*, as can be observed in the description of this new species which, characteristically, shows marked violet colours on the stipe of young specimens, and moreover, its pileus is not rimose at the marginal zone and often splits into small, concentric, appressed scales when extended, the spores lack a supra-hilar depression, without conical-pointed apex and the cystidia show a somewhat thicker wall ($W_m = 1.8$).

It is noteworthy that both taxa occur in locations with a certain human influence, in ruderalised areas and nitrified soils (parks, gardens). *Inocybe paleoveneta* has been found under *Betula* and *Carpinus* (*Betulaceae*) in Italy, but other collections from Germany, originally determined as *I. grammopodia*, grew in garden areas under *Quercus* and *Tilia* (Table 1). This ruderal habitat is also common for *I. pusio* (Vauras, pers. comm.; LUDWIG 2017) and the new species *I. ianthinopes* Pancorbo, G. Muñoz & Esteve-Rav. (described below). The lectotype specimen

of *I. pusio*, designated by BANDINI & al. (2021), shows a different composition of the stipitipellis and, particularly, caulocystidia which extend up to 1/4–1/3 of the stipe apex; other differences are the regularly broad and ventricose hymenial cystidia and the absence of a developed veil in early stages.

Inocybe ianthinopes Pancorbo, G. Muñoz & Esteve-Rav., *sp. nov.*

MycoBank: MB 841754

Holotypus: SPAIN. **Zaragoza:** Zaragoza, Parque Grande José Antonio Labordeta, Jardín Botánico, 41.633972 -0.895.444, 243 m, on muddy soil, under *Tilia tomentosa*, *Prunus laurocerasus*, *Arbutus unedo* and *Berberis vulgaris*, 25-X-2012, leg. G. Muñoz, AH 40370 (ITS GenBank OL352086). *Isotypus* in GM1589.

Figs. 4-5

Etymology: *ianthinopes* (Latin), from *ianthinum* = violaceous and *pes* = stipe, a typical feature of this species with intense violet stipe.

Diagnosis: morphologically characterised by the slender habit, fibrillose to woolly pileus when young and sublageniform cystidia with long protruding neck; *I. knautiana* is similar macroscopically but differs in differently shaped cystidia. The ITS differs in BLAST searches from similar species like *I. pusio*, *I. derbschii* (89% of similarity, KJ432287 —sub *I. pusio*—, MW845868 —sub *I. derbschii*—) and *I. caesaraugustae* (88.5% of similarity, MW354973 —sub *I. aff. grammopodia*).

Basidiomata agaricoid and stipitate. **Pileus** 10–20(–25) mm diam, at first conical, campanulate to conico-convex, then plano-convex, umbonate in adult specimens, umbo either distinctly convex to shallow, not hygrophanous; margin hardly involute to straight when young, regular to wavy in old specimens, velipellis not distinct; colour uniformly chestnut brown to dark brown (Mu 5YR 4/3–6, 7.5YR 4/3–4), in AH 48228 paler to grey–yellow brown (10YR 3/4–6); surface dry, finely tomentose to felty lanose, especially at the centre, sometimes radially virgate, fibrillo-squamulose in adult specimens towards the margin, then showing a paler background, extreme margin paler in young specimens by the presence of remnants of a fragile whitish cortina. **Lamellae** subdistant to moderately crowded [$L = 28$ –34; $l = 1$ (–2)], emarginate, ventricose, typically lilac to violet lilaceous when young, then glaucous to argillaceous to brown argillaceous, finally tobacco-brown, edge white, finely fimbriate. **Stipe** 20–40 × 2–3(–4) mm, fibrose, frequently curved or sinuose, cylindrical to progressively enlarged towards the base; colour intensely and uniformly violaceous to bluish violet at first, becoming lilac to lilac ochraceous, extreme base

Fig. 4. *Inocybe ianthinopes*. A: Collection AH 40370. Holotype. B: Collection AH 48228. Scale bar: 20 mm.

Fig. 5. *Inocybe ianthinopes* AH 40370. Holotype. A: Basidiospores. B, D: Cheilocystidia. C: Pleurocystidia. E: Caulocystidia and caulocystidioid hairs. Scale bar: A: 10 μm , B-D: 50 μm . Mounting media: A-E: $\text{NH}_3(\text{aq})$.

often paler and even whitish with age; surface typically covered by abundant appressed white fibrils, often mimicking the violaceous underground when young, persistent to ephemeral, fibrillose to falsely floconose at the extreme apex. *Cortina* white, present, fleeting. *Context* fibrose, whitish at the pileus, lilac or pale violaceous at the stipe cortical zone and stipe apex, paler towards the inner flesh. *Smell* strongly spermatic, taste not significant.

Basidiospores smooth, brownish, amygdaliform with ogival apex, frequently papillate, some with a suprahilar depression $(7.9\text{--}8.4\text{--}10.2\text{--}11.1) \times (4.4\text{--}4.8\text{--}5.7\text{--}6.5)$. $Q = (1.4\text{--}1.6\text{--}2\text{--}2.3)$, ($L_m = 9.3\text{ }\mu\text{m}$, $W_m = 5.2\text{ }\mu\text{m}$, $Q_m = 1.8$; $n = 163$, $N = 2$), slightly thick walled ($-0.5\text{ }\mu\text{m}$), with a central large vacuole. *Basidia* hyaline, claviform, 4-spored, $(23.7\text{--}25.3\text{--}33\text{--}37) \times (7.3\text{--}7.7\text{--}9.4\text{--}11.6)\text{ }\mu\text{m}$, ($L_m = 29.2\text{ }\mu\text{m}$, $W_m = 8.6\text{ }\mu\text{m}$; $n = 50$, $N = 2$). Gill edge nearly homogeneous and

sterile in young specimens. *Cheilocystidia* (33-) 40-57 (-65) $\times$ (9.7-)11-16(-23) μm , ($L_m=50$, $W_m=13$; $n=86$, $N=2$), very abundant, hyaline or showing a pale yellowish intracellular pigment, very crystalliferous at apex, morphologically variable, often narrowly lageniform to ventricose-lageniform with a distinct, often undate neck, less frequently subcylindrical to subfusiform, thin walled [0.5-1.0(-1.5) μm], with hardly any reaction to ammonia or very pale yellow; *paracystidia* abundant by zones, clavate to pyriform, hyaline, sometimes septate, 15-35 $\times$ 8-17(-20) μm . *Pleurocystidia* similar and morphologically variable like cheilocystidia, usually longer, (42-)46-65(-70) $\times$ (9.8-)12-17(-22) μm , ($L_m=55$, $W_m=14$, $n=86$, $N=2$). *Stipitipellis* a cutis of parallel hyphae of 2.5-6.6 μm wide. *Caulocystidia* very rare, similar to hymenial cystidia, only present at the extreme apex (1/10) or even absent in some collections; in AH 48228 somewhat more frequent, claviform to subcylindrical, some metuloid and crystalliferous, 30-59 $\times$ 10-17 μm ; stipe apex provided with numerous hyaline cylindrical to subclavate hairs with rounded apex (fibrillose), narrow, 5-12 μm wide. *Clamps* present.

Habitat & Distribution: This species also seems to have similar ecological preferences as *I. pusio*. The holotype was found very close to and in the same habitat as *I. caesaraugustae*, which leads us to assume its preference for ruderalised and somewhat nitrified areas like parks and gardens, especially next to broad-leaved trees and probably also under conifers. The other collection found in the Iberian Peninsula, however, was collected in a damp hazel spot in the vicinity of a *Pinus sylvestris* forest, on calcareous soil and poorly human-influenced. Similar sequences from GenBank that correspond to *I. ianthinopes* (Tab. 1) indicate a wide distribution in Europe, in wet and temperate areas (Germany, Italy, Sweden—Larsson pers. comm.) in well drained soils. Thus, it is very likely that some collections identified as *I. pusio* correspond to this species due to their great macroscopic similarity; this is the case of both collections cited by BANDINI & al. (2019), referred to *I. cf. pusio*.

Additional material studied: SPAIN. **Huesca:** Seira, barranco de Barbaruens, 42.515544 0.376183 (± 100 m), 1404 m, on calcareous humid soil under *Corylus avellana*, 28-VIII-2018, leg. F. Pancorbo, AH 48228, (ITS GenBank OL352087). Duplicate in FP18092803.

Comments: *Inocybe ianthinopes* is a small and slender species, characterised by a brown to chestnut-brown pileus, with a woolly-felted appearance, sometimes tabby to felted-reticulate at maturity (its appearance is often reminiscent of species of the *I.*

cinnamata group) and the stipe is intensely violet in young stages, often covered by numerous whitish veil fibrils that disappear with age. The cystidia are often lageniform to subcylindrical, sometimes ventricose, but have always a marked tendency to attenuate into a distinctive protruding neck, the walls being often undate and typically very narrow, not yellowing in ammonia solutions. Caulocystidia are either absent or present in the stipe insertion zone. The appearance of the pileus and cystidia seem to be characteristic of this species and facilitate its macroscopic and microscopic separation from phylogenetically close taxa such as *I. pusio*, *I. paleoveneta* and *I. caesaraugustae*, which have a smoother and more radially fibrous pileus, at least when young. *Inocybe knautiana* Bandini & B. Oertel is sometimes reminiscent of *I. ianthinopes* because of the felted appearance of the pileus and the colour of the stipe; however, *I. knautiana* occupies an isolated phylogenetic position and shows broad, fusiform hymenial cystidia (BANDINI & al. 2021).

Our analyses (Fig. 1) show that two sequences (MH366588 and MH366589) from specimens published by BANDINI & al. (2019) as *I. cf. pusio*, and another one (JF908164), as *I. amethystina*, belong in fact to *I. ianthinopes*. However, our concept of *I. pusio*, based on the study of lectotype, which coincides with that of J. Vauras and partially with that of KUYPER (1986), corresponds to a species with caulocystidia in the upper third of the stipe, with radially fibrous pileus, smooth in appearance, barely somewhat felted or tomentose in the centre when young, with not or slightly developed veil around disc; typically the hymenial cystidia are regularly broad (>20 μm in the lectotype), ventricose to broadly fusiform, with walls 1.5-2 μm thick, hyaline or hardly yellowish in ammonia.

***Inocybe velatipusio* Esteve-Raventós & Pancorbo, sp. nov.**

MycoBank: MB 841755

Holotypus: FRANCE. **Nouvelle-Aquitaine:** Pyrénées-Atlantiques, Osse-en-Aspe, Forêt d'Issaux, 43.00250 -0.71139, 938 m, under *Corylus avellana* in a *Fagus sylvatica* and *Abies alba* forest, 11-X-2016. leg. M.Á. Ribes, J. Cuesta, J. F. Mateo, F. Pancorbo, AH 47725, (ITS GenBank OL352088). **Isotypus** in FP16101104.

Figs. 6-7

Etymology: *velati*, from Latin *velatus* = provided with a veil, and *pusio*, in reference to *Inocybe pusio*, which is apparently a close species.

Diagnosis: *Inocybe velatipusio* is characterised by the smooth pileus covered by a well-developed velipellis, at least in young stages, and the deep violet stipe, devoid of caulocystidia or only present in the

Fig. 6. *Inocybe velatipusio*. A. Collection AH 47725. Holotype. B. Collection AH 45163. C. Collection AH 25337 (collection with no visible velipellis). D. Collection AH 26788 (collection with no visible velipellis). Scale bar: 10 mm = A. Photo C: A. Rocabrana

insertion zone. With the exception of the stipe colour, it morphologically resembles *I. griseovelata* Kühner due to the conspicuous presence of the velipellis. It differs from *I. pusio* by the different composition of the stipitipellis and the absent or rudimentary veil in the latter. Phylogenetically it is not close to any species with violet colours (only *I. dryadiana* Bandini & B. Oertel blasts at ITS 83%); with a different stipe colour, devoid of violaceous tones, the closest species at ITS are *I. grusiana* Bandini & B. Oertel (89%), *I. virgatula* Kühner and *I. fuscidula* Velen. s. auct. non Bandini (84%).

Basidiomata agaricoid and stipitate. *Pileus* 20-25(-32) mm diam, not hygrophanous, at first hemisphaerical to conico-convex, then plano-convex, umbo usually distinctive, broadly conical or shallow, rarely absent; margin slightly incurved when young, finally straight, regular to wavy in old and expanded specimens; velipellis distinct whitish to pale greyish, usually persistent at the centre, but ephemeral in

washed or aged collections; colour uniformly chestnut brown to grey brown (Mu 5YR 4/3-6, 7.5YR 4/3-4) or even purplish brown in embedded specimens (AH 45163); surface dry, smooth, radially fibrose, finely rimose towards the margin with age, taking on a “virgate” aspect, extreme margin sometimes showing the presence of remnants of a cobwebby whitish cortina. *Lamellae* moderately crowded [$L = 30-40$; $l = 1(-2)$], emarginate, ventricose, whitish, glaucous, often tinged pale lilaceous when young, then argillaceous to brown argillaceous, finally tobacco-brown; edge whitish or paler, finely fimbriate. *Stipe* 20-40 (-50) $\times$ (3-)4-6(-7) mm, fibrose, frequently curved or sinuose, cylindrical to progressively enlarged towards the base; colour intensely and uniformly violaceous to pinkish violaceous at first, becoming lilac to lilac ochraceous, base sometimes paler and even whitish with age; surface fibrillose, covered by abundant appressed white fibrils, normally persistent, fibrillose to falsely fibro-floconose at the extreme apex or upper third. *Cortina* whitish, fugacious. *Context* fibrose,

Fig. 7. *Inocybe velatipusio* AH 47725. Holotype. A. Basidiospores. B. Cheilocystidia C. Pleurocystidia D. Caulocystidia at apex of stipe. Scale bar: 10 µm = A, 50 µm = B-D. Mounting media µm: $\text{NH}_{3(\text{aq})}$ =A-D

whitish at the pileus, pale lilac at the cortex and upper part of the stipe, paler to whitish towards the inner flesh. *Smell* subspermatic, taste not significant.

Basidiospores smooth, brownish, amygdaliform with ogival apex, in some collection often sub-papillate, some with a suprahilar depression $(8\text{--}9.1\text{--}12.4\text{--}(15) \times (4.9\text{--}5.2\text{--}6.8\text{--}(7.7)) \mu\text{m}$, $Q = (1.4\text{--}1.6\text{--}1.9\text{--}(2.3))$ ($L_m = 10.5 \mu\text{m}$, $W_m = 5.9 \mu\text{m}$, $Q_m = 1.8$; $n = 201$, $N = 7$), slightly thick walled ($-0.5 \mu\text{m}$), with a central large vacuole. *Basidia* hyaline, claviform, mostly 4-spored, $(20\text{--}24\text{--}39\text{--}(41)) \times (6.7\text{--}7.9\text{--}12\text{--}(13)) \mu\text{m}$ ($L_m = 31 \mu\text{m}$, $W_m = 9.6 \mu\text{m}$; $n = 82$, $N = 2$). Gill edge nearly homogeneous and sterile. *Cheilocystidia* $(35\text{--}48\text{--}70\text{--}(83)) \times (9.6\text{--}13\text{--}21\text{--}(25)) \mu\text{m}$ ($L_m = 59 \mu\text{m}$, $W_m = 17 \mu\text{m}$, $n = 109$, $N = 7$), (in AH 48163 reaching $117 \mu\text{m}$), very abundant, hyaline or showing an ochraceous intracellular pigment, crystalliferous at apex (sometimes showing the apex covered by microcrystals), morphologically variable, often fusiform to subcylindrical (in some occasions subcapitate), sometimes sublageniform or claviform, thin to medium thick-walled $(1.1\text{--}1.3\text{--}2.2\text{--}(2.5)) \mu\text{m}$, with very pale yellow reaction to ammonia; *paracystidia* numerous in some cases, clavate to subcylindrical, hyaline, sometimes septate. *Pleurocystidia* numerous, similar and morphologically variable like cheilocystidia, $(39\text{--}53\text{--}80\text{--}(92)) \times$

$(11\text{--}14\text{--}21\text{--}(27)) \mu\text{m}$, ($L_m = 64 \mu\text{m}$, $W_m = 18 \mu\text{m}$, $n = 116$, $N = 7$), hyaline or showing an ochraceous intracellular pigment. *Stipitipellis* a cutis of parallel hyphae of $2.5\text{--}6.3 \mu\text{m}$ wide. *Caulocystidia* sublageniform to subcylindrical $(46\text{--}48\text{--}68\text{--}(92)) \times (14\text{--}15\text{--}19\text{--}(20)) \mu\text{m}$, ($L_m = 59 \mu\text{m}$, $W_m = 17 \mu\text{m}$, Holotype), present in the apical part of the stipe (in upper 1/10), mixed with numerous hyaline cylindrical to subclavate hairs with rounded apex (fibrillose), $4\text{--}11 \mu\text{m}$ wide. *Clamps* present.

Habitat & Distribution: *Inocybe velatipusio* seems to show a preference for *Fagaceae* forests, especially in western Mediterranean and atlantic-submediterranean areas (as in the holotypic locality), on preferably acidic soils. We have found it in both evergreen *Quercus ilex* and deciduous forests, such as *Q. faginea* and *Fagus sylvatica* at higher elevation. For the moment we only know it to be present in Spain, southern France as well as in the eastern mediterranean beech and oak forests in Iran (Hyrcanian forests), judging from a GenBank sequence (FR852257, Table 1).

Additional material studied: SPAIN. Jaén: Palomares, in *Quercus ilex* forest, 17-V-1996, leg. F. Jiménez, AH 19399, duplicate in Herb. F. Jiménez.

Barcelona: Premià de Dalt, camino a Sant Bertomeu de Cabanyes, mixed forest of *Pinus pinea* and *Quercus ilex*, 22-X-1999, leg. A. Rocabruna, AH 25337. **Madrid:** Rozas de Puerto Real, embalse de Los Morales, 780 m, under *Quercus pyrenaica* in acidic soil, 14-V-2000, leg. F. Esteve-Raventós & M. Villareal, AH 26788. **Ciudad Real:** Viso del Marqués, Arroyo de La Poveda, *Quercus faginea* forest, in acidic soil, 29-V-2004, leg. F.D. Calonge, AH 34397, (ITS GenBank OL352089). **Toledo:** Cantera de San Pedro, Montesclaros, 40.087844 -4.934242, 485 m, in *Quercus faginea* forest with *Cistus monspeliensis* in acidic soil, 27-XII-2014, leg. F. Pancorbo, AH 45163, duplicate in FP14122713 (ITS GenBank OL352090).

Comments: *Inocybe velatipusio* reminds of *I. pusio* provided with an extensive development of the universal veil or velipellis, although in some collections this can be ephemeral, probably due to environmental conditions (Fig. 6, C-D). This veil is also conspicuous along the stipe, forming a thin whitish to greyish mantle of hyphae, sometimes broken into fine apparent strands. From the collections studied and sequences in GenBank, the geographical distribution of *I. velatipusio* seems to be limited to southern Europe, in temperate or warm climates, extending as far as Iran, always linked to *Fagaceae*.

In our analyses of the ITS region (Fig. 1), *I. velatipusio* does not nest in the clade comprising the rest of the species of the *I. pusio* group, and thus, it might be that it is distant to them. BLAST searches show a similarity of 83.4% with *I. fuscidula* Velen. s. Kuyper (GenBank MW856438 and MT005928), non Bandini or *I. virgatula* Kühner. Interestingly, the appearance of the pileus of *I. velatipusio* is very reminiscent of these species, appearing smooth and radially fibrous, although this character is often mimicked by the apparent whitish to pale grey veil. Another particular character of *I. velatipusio* is the presence of caulocystidia restricted to the insertion zone of the stipe (upper 1/10), although in some collections they may appear absent; the caulocystidia are elongate sublageniform to subcylindrical (Fig. 7), and appear clustered with transitional forms to elongate elements (hairs), but we have never observed the presence of claviform or spheropedunculate paracystidia. Elongate and narrow caulocystidia are also reported for *I. derbschii* Schwöbel & Stangl (BANDINI & al. 2019), *I. grusiana* Bandini & B. Oertel, *I. ghibliana* Bandini & B. Oertel and *I. plurabellae* Bandini, B. Oertel & U. Eberh. (BANDINI & al. 2021). However, all these species lack lilac or violet shades on the stipe and are part of the “complex” *I. virgatula/fuscidula* s. auct. pl.”.

Inocybe griseovelata also shows similarities with *I. velatipusio* in relation to the presence of a pale greyish veil on the pileus. KÜHNER (1955) pointed out

a stipe with some pinkish shades in the apical zone, the rest being whitish. A specimen studied by us in 1988, marked by Poirier in 1985 as lectotype—but never proposed as such in an effective publication—, lacks any stipe fragment, so that the distribution of the caulocystidia could not be checked. Recently, BANDINI & al. (2021) mentioned that this specimen (through a communication by P. Clerc, curator at G) is lost. Hence, a specimen collected in the same locality, collected in a different year, was designated as lectotype by BANDINI & al. (2021), but sequence data from it are unavailable. *Inocybe dryadiana* also shows violaceous colourations on the stipe, but the colour of the squamulose pileus is tawny brown, it does not show a developed velipellis and the cystidia are different in size and shape, while *I. grusiana* also shows a very evident velipellis but lacks violaceous or lilac colours (BANDINI & al. 2021).

The morphological study of an isotype of *Inocybe pusio* f. *velata* Reumaux (Herb. P. Reumaux n° 500, holotype in PC 99597) probably indicates that this taxon could be related or co-specific with *I. velatipusio*; the smooth, radially fibrous appearance of the pileus, the brown colour of the pileus and the presence of an apparent veil are shared characters. This similarity can also be established for microscopic characters (see REUMAUX 1982; MOËNNE-LOC-COZ & al. 1988). However, our attempts of obtaining genetic material proved unsuccessful and refrains us from proposing their conspecificity. Although the type material of *I. pusio* f. *velata* was reported under *Populus* on calcareous ground, the Latin diagnosis specifies “...*Populis tremulis Quercetisque*...”, which could support the hypothesis of its relationship with *Fagaceae*.

Inocybe pusio, as previously stated (see remarks on *I. caesaraugustae*), shows caulocystidia on at least the upper 1/4-1/3 of the stipe and no apparent velipellis on the pileus.

New combinations:

Inocybe floccipes (Esteve-Rav. & Fouchier) Esteve-Rav. & Bizio, **comb. & stat. nov.**

Mycobank: MB 841756

Basionym: *Inocybe pusio* var. *floccipes* Esteve-Rav. & Fouchier, *Cryptog. Mycol.* 25(2): 108. 2004. Mycobank MB 487652

Fig. 8

Additional material studied: FRANCE. **Alpes de Haute-Provence:** Méailles, 1400 m, under *Castanea sativa*, 22-X-1998, leg. F. Fouchier, F. Fouchier n° 98085, duplicate in M. Bon 98157 (LIP). ITALY. **Apulia:** Brindisi, Bosco Colemi Tutturano, under *Quercus*, 22-IV-2009, leg. C. Agnello, Herb. E. Bizio,

Fig. 8. *Inocybe floccipes*. A. Collection AH 40368, La Rioja, Spain. B. Collection 20140601, Pregarje, Slovenia. C. Collection AH 46767, Mallorca, Spain. Photo B: E. Bizio

duplicate in AH 40051 (ITS GenBank OL352061). **PORTUGAL. Beira Litoral:** Leiria, Mata de Curvachia, 39.713632 -8.766669, 75 m, under *Quercus faginea*, 9-XI-2000, leg. F. Esteve-Raventós, AH 29885 (ITS GenBank OL352059). **SPAIN. Barcelona:** Sant Celoni, Rectoria d'Olzinelles, 41.665764 -2.512440 (± 200 m), 250 m, in *Quercus ilex* subsp. *ilex* forest with *Corylus avellana*, leg. F. Esteve-Raventós, G. Moreno, J. Llistosella & A. Rocabruna, AH 24954. **Burgos:** Valle de Mena, Santuario de Cantonad, 43.082158 -3.356266, 500 m, in *Fagus sylvatica* forest, 14-X-1998, leg. F. Esteve-Raventós & J.M. Barrasa, AH 24988 (ITS GenBank OL352058). **Islas**

Baleares: Mallorca, Campanet, Fonts Ufanes, 75 m, in *Quercus ilex* subsp. *ilex* forest, 22-XI-2012, leg. F. Esteve-Raventós, J. L. Siquier & J. C. Salom, AH 46767 (ITS GenBank OL352060). **La Rioja:** Tudellilla, in *Quercus ilex* subsp. *ballota* forest, 8-XI-2008, leg. G. Muñoz, AH 40368, duplicate in GM1428. **Navarra:** Ultzama, Lizaso, Club de golf Ultzama, 520 m, under *Quercus robur*, 14-VI-2011, leg. A. Caballero, AH 40327, duplicate in AC4712. **Taragona:** Mas de Barberans, barranco de Retaule, 40.744535 0.264012, 1000 m, in *Fagus sylvatica* forest, 16-X-2001, leg. F. Esteve-Raventós, AH 30626

(Holotype) (ITS GenBank OK414030, LSU GenBank OK422243).

Comments: Our molecular analyses (including sequences obtained from the holotype), leave no doubt that this is a species distinct from those belonging to the “pusio” group (see Fig. 1 *Inocybe* cf. *pusio*). *Inocybe floccipes* was described by ESTEVE-RAVENTÓS & FOUCHIER (2004), on the basis of an invalid description by Fouchier in BON (1999) of “*I. pusio* f. *laticaulocystidiata*”, and characterised by the presence of abundant caulocystidia that may even reach the base of the stipe, or at least present at the lower 2/3 of it. The cortina is extremely fleeting, and could only be observed in one of the paratypes on a small basidioma, still at a very young stage. There are no remains of veil on the pileus in any of the collections studied (see Fig. 8).

Since the original description, this taxon has been the subject of some subsequent studies, such as that by CABALLERO & MUÑOZ (2011), who have found it in La Rioja (Spain) several times under holm oaks (*Quercus ilex* subsp. *ballota*). Those authors pointed out to the presence of a very fleeting cortina, which supports our observations and also a relationship with the *I. pusio* group and not with *Inocybe* sect. *Splendentes* R. Heim (see Fig. 1). CARTERET & REUMAUX (2011) described *I. floccipes* also it from material collected in France under *Fagaceae* (*Fagus* and *Castanea*).

Judging from some collections made by our colleague Enrico Bizio (Italy), it appears that *I. floccipes* is widespread in the southern areas of Europe. It is known from France, Italy, Portugal, Slovenia (GenBank MW027928, MW027944, MW 027970, MW027991) and Spain, in warm or temperate climates. *Inocybe floccipes* is clearly associated with *Fagaceae* (*Quercus*, *Fagus*, *Castanea*), often on acidic soils, less often on calcareous substrate.

Inocybe paleoveneta (Bizio & A. Castellan) Esteve-Rav. & Pancorbo, **comb. & stat. nov.**

MycoBank: MB 841757

Basionym: *Inocybe grammopodia* var. *paleoveneta* Bizio & A. Castellan, *Micol. Veg. Medit.* 32(2): 112. 2018. MycoBank MB 824287.

Figs. 9-10

Material studied: SPAIN. **Madrid:** Humera, aprox. 650 m, in a private garden under *Carpinus betulus*, 15-IX-2021, leg. G. Moreno, AH 51448 (ITS GenBank OL583808).

Additional material studied: *Inocybe grammopodia* Malençon: MOROCCO. **Moyen Atlas:** Azrou, plateau de Scheb, Ich Ouharrock, 1950 m, under *Cedrus*, 11-V-1952, leg. G. Malençon, Herb. Malençon n° 2158 (LIP, holotype); *Ibidem*, 25-IV-1952, leg. G. Malençon, Herb. Malençon n° 2107 (LIP).

Comments: The study made of the specimens collected in North Africa, including the holotype of *I. grammopodia* (MPU), fits perfectly with the original detailed description (MALENÇON & BERTAULT 1970). The protologue mentioned the characters of the stipe, noted as “...*bulbous submarginate, ex albo roseo-brunneo vel pallide fulvescente, toto pruinoso,...*”, characters which do not fit the description of *I. paleoveneta*, characterised by the pinkish-lilac tones of the stipe and, above all, by the presence of caulocystidia only at the upper end of the stipe. In our study of the Malençon material, we confirmed the presence, sometimes numerous, of caulocystidia in the lower third of the stipe, what was already observed by KUYPER (1986), who previously revised the type collection (Fig. 11). The presence of a cortina is not mentioned in the original description. It is quite possible that *I. grammopodia* and *I. paleoveneta*, that sometimes show a longitudinally striped stipe, could be related to each other. On the other hand, collections of *I. grammopodia* do not show the pink-lilac tones typical in *I. paleoveneta* and represented on all photographs in BIZIO & CASTELLAN (2018), but the stipe, which is originally whitish, turns slowly fuscous to reddish brown with age. Other additional differences such as the habitat and distribution also suggest the consideration of *I. paleoveneta* as a separate species.

We have also studied numerous collections of *I. grammopodia* from the Iberian Peninsula, that fit very well with the original diagnosis, always in Mediterranean environments with conifers and broad-leaved trees (Fig. 11). In addition, *I. grammopodia* is microscopically characterised by its narrow spores often showing a typical supra-hilar depression that makes the spores appear sub-phaseoliform ($Q_m=1.95$ in the holotype collection, $n=30$) (Fig. 11). For all these reasons, we disagree with BIZIO & CASTELLAN (2018) and BANDINI & *al.* (2019) who considered *I. paleoveneta* a variety of *I. grammopodia*. Regrettably, the holotype of *I. grammopodia* has not been sequenced, but we think that it will prove unsuccessful as the vouchers were treated with Cl_3Hg vapours for preservation.

Discussion

The recent use of molecular tools for taxonomic purposes has reliably clarified the polyphyletism of the traditional *I. sect. Lilacinae* (JACOBSON 2008; JACOBSON & LARSSON 2012), typified by *I. pusio* (BON 1977). Based mostly on genetic studies, supported by further morphological and ecological characters, we present in this contribution some taxonomic novelties of species reminiscent in appearance and habit of *I. pusio*. Morphological diagnostic

Fig. 9. *Inocybe paleoveneta* AH 51448. A. Basidiomata. Scale bar: 20 mm = A

Fig. 10. *Inocybe paleoveneta* AH 51448. A. Basidiospores. B. Pleurocystidia C. Cheilocystidia D. Caulocystidia at apex of stipe. Scale bar: 10 μm = A, 50 μm = B-D. Mounting media: $\text{NH}_{3(\text{aq})}$ = A-D

Fig. 11. *Inocybe grammopodia*. A. Collection AH 46786. B. Collection AH 25042. C. Voucher, original drawings and notes taken by T.W. Kuyper on the holotype of *I. grammopodia*. D. SEM Spores from Holotype Herb. Malençon n° 2158 (LIP)

characters are, however, rather subtle. Through the interpretation of our analyses, it is obvious that the distribution of caulocystidia and the aspect of the pileus are characters to be taken into account in the recognition of some species reminiscent of *I. pusio*. Some of them (e.g. *I. velatipusio* Esteve-Rav. & Pancorbo) belong to other clades or phylogenetic groups despite their morphological resemblance. In addition, colours should be interpreted with caution, as in many species the purplish or lilac pigments tend to fade, to go unnoticed or even disappear, especially with age, dehydration or in dried herbarium material. In dried collections these pigments often take on ochraceous or yellowish tones or colourations that make a correct determination difficult. It is therefore very important to have fresh samples in good condition and, if possible, specimens in different development stages, in order to recognise these possible chromatic changes.

The study of type collections is nowadays a real necessity to be able to corroborate the original interpretations. In this case, the morphological study of the original material of *I. pusio* (Herb. P. Karsten n° 1608a, in TUR-A as H-6006856) has been essential for the correct taxonomic framing of the species presented here and their possible confusions. This study was carried out by us in 1999, and our observations match those of the revision

and study made by J. Vauras some years earlier (see ESTEVE-RAVENTÓS & FOUCHIER 2004); the presence of caulocystidia in the upper 1/3-1/4 of the stipe and the ventricose hymenial cystidia are particular to *I. pusio*. Three collections determined as *I. pusio* by J. Vauras from Finland, Estonia and Sweden were sent for our study from TUR-A; all of them were sequenced and deposited in GenBank under accession numbers KJ432286, KJ432287 and KJ432288, and conform to the sequence under accession number JF908241 in GenBank (as *I. amethystina* Kuyper). This set of sequences might represent Karsten's species (see Table 1), pending that barcode sequences are obtained from the lectotype designated by BANDINI & *al.* (2021) (coll. H-6006856, no. 1608a, MBT10000534) which corresponds to the same collection studied by us in 1999. For this reason, awaiting the ongoing molecular study by our colleague E. Larsson (University of Gothenburg, Sweden), we refrain at the moment from adopting this clade as unambiguously corresponding to the type of P. Karsten, and refer to it as *I. cf. pusio*. Only if sequencing of the lectotype proved to be unsuccessful, any of the collections authenticated by J. Vauras from Finland would be good candidates for proposing an epitype.

Artificial key to the species treated in this work

All the species treated in this contribution share a more or less intense lilac or violet colours in the stipe and smooth spores, and are thus macroscopically reminiscent of *I. pusio*.

- 1.-Caulocystidia absent or only present at the extreme apex or insertion of the stipe; cystidia walls narrow up to 1.5 µm thick.....2
 - 1.-Caulocystidia present at least at the upper 1/3-1/4; cystidial walls 1.5-2.5 µm thick.....5
 - 2.-Spores ellipsoid to subamygdaliform with not distinctly pointed or papillate apex
..... *I. caesaraugustae*
 - 2.- Spores amygdaliform with pointed apex, sometimes subpapillate3
 - 3.-Pileus felted to woolly-fibrillose, chestnut brown to brownish without a developed velipellis. In forests, parks and gardens *I. ianthinopes*
 - 3.-Pileus smooth, radially-fibrillose, often covered with a whitish velipellis, especially in young stages4
 - 4.-Under various broadleaved trees in parks and gardens *I. paleoveneta*
 - 4.-Under *Fagaceae* (*Quercus*, *Fagus*, *Castanea*) .
..... *I. velatipusio*
 - 5.- Caulocystidia present in upper 1/3-1/4 of stipe; hymenial cystidia typically ventricose. In ruderalised areas, parks and gardens, under broadleaved trees, not necessarily *Fagaceae* *I. pusio*
 - 5.-Caulocystidia abundant, reaching the lower 2/3 of the stipe; hymenial cystidia mostly fusiform; under *Fagaceae* *I. floccipes*

Acknowledgements

Fernando Esteve-Raventós and partners were financially supported by FEDER/Ministerio de Ciencia, Innovación y Universidades – Agencia Estatal de Investigación (Spain)/Project CGL2017-86540-P, in which this contribution is included.

We would like to thank Profs. Alberto Altés and Gabriel Moreno (Universidad de Alcalá,), for kindly authorizing us to publish the SEM photograph of the spores of the holotype of *I. grammopodia* and also the macroscopic photograph of *I. paleoveneta*. To our friend the late Agustín Caballero, for providing us with samples and information; to Enrico Bizio for sharing his data on *I. floccipes*; to Caroline Loup (Curator of the MPU Herbarium, Montpellier) for her kindness and facilities for the study of G. Malençon Herbarium; to Jukka Vauras for the information provided and the sending of samples for study and sequencing. To Lidia Plaza for generating

some of the sequences included in this work. Finally, Yolanda Turégano would like to thank Miguel Blázquez and Paula Garrido for his methodological and graphic advice regarding phylogenetic analyses and edition of the tree figure, respectively. We also thank J. L. Baptista-Ferreira and G. Marques for translating the summary to portuguese.

Our thanks to E. Larsson and P. B. Matheny for reviewing our manuscript and providing us with kind suggestions that implied the scientific improvement of this work.

References

- ALTSCHUL SF, GISH W, MILLER W, MYERS EW & LIPMAN DJ (1990) “Basic local alignment search tool.” *J. Mol. Biol.* 215: 403-410.
- BANDINI D, OERTEL B, PLOCH S, ALI T, VAURAS J, SCHNEIDER A, SCHOLLER M, EBERHARDT U & al. (2019) Revision of some central European species of *Inocybe* (Fr.: Fr.) Fr. subgenus *Inocybe*, with the description of five new species. *Mycol. Progress* 18: 247-294. DOI: <https://doi.org/10.1007/s11557-018-1439-9>
- BANDINI D, OERTEL B & EBERHARDT U (2021) A fresh outlook on the smooth-spored species of *Inocybe*: type studies and 18 new species. *Mycol. Progress* 20: 1019-1114. DOI: <https://doi.org/10.1007/s11557-021-01712-w>
- BIZIO E & CASTELLAN A (2018) *Inocybe acuto-fulva* e *Inocybe grammopodia* var. *paleoveneta*, due nuovi taxa dall’Alta Marca Trevigiana (Trevviso, Veneto, Italia). *Micol. Veg. Medit.* 32(2): 103-124.
- BON M (1997) Clé monographique du genre *Inocybe* (Fr.) Fr. (2ème partie: Sous genre *Inocybe*=*Inocybium* (Earle) Sing.). *Doc. Mycol.* 27 (108): 1-77.
- BON M (1999) Étude de quelques *Inocybes* méridionaux aux Journées de la F.A.M.M. à Annot (Octobre 1998). *Bull. Semestr. Féd. Assoc. Mycol. Médit.* 15(1): 3-12.
- CABALLERO A & MUÑOZ G (2011) Algunas especies raras o interesantes de Agaricales recolectadas en la Península ibérica. *Bol. Micol. FAMCAL* 6: 39-51.
- CARTERET X & REUMAUX P (2011) Miettes sur les *inocybes* (6^e série). Étude de quelque nains des feuillus de la plaine, accompagnée d’une clé de détermination des taxons de la section *Lilacinae* R. Heim. *Bull. Soc. Mycol. France* 127 (1-2): 1-53.
- CZISMÁR M, CSEH P, DIMA B, ORLÓCI L & BRATEK Z (2021) Macrofungi of urban Tilia avenues and gardens in Hungary. *Glob.*

- Ecol. Conserv.* 28 e01672. DOI: <https://doi.org/10.1016/j.gecco.2021.e01672>
- ESTEVE-RAVENTÓS F & FOUCHIER F (2004) *Inocybe pusio* var. *floccipes* var. nov. and some observations on the variability in this species. *Cryptog. Mycol.* 25(2): 107-114.
- HEIM R (1931) *Le genre Inocybe. Encyclopédie Mycologique* Vol. I. Paris, P. Lechevalier, 431 p.+ 35 pl.
- JACOBSON S (2008) *Inocybe*: 868-906. In: KNUDSEN H. & VESTERHOLT J. (ed.). *Funga Nordica*. Nordsvamp. Copenhagen.
- JACOBSON S & LARSSON E (2012) *Inocybe*: 981-1021. In: KNUDSEN H. & VESTERHOLT J. (ed.). *Funga Nordica* (2nd edition). Nordsvamp. Copenhagen.
- KÜHNER R (1955) Compléments à la “Flore Analytique” V) *Inocybe* léiosporés cystidiés. Espèces nouvelles ou critiques. *Bull. Soc. Nat. Oyonnax* 9 (Suppl.): 3-95.
- KÜHNER R & ROMAGNESI H (1953) *Flore analytique des champignons supérieurs (Agarics, Bolets, Chanterelles) comprenant les espèces de l'Europe occidentale et centrale*. Masson et Cie. Paris. 556 pp.
- LARSSON A (2014) AliView: a fast and lightweight alignment viewer and editor for large data sets. *Bioinformatics* 30(22): 3272-3276. DOI: <https://doi.org/10.1093/bioinformatics/btu531>
- KUYPER TW (1986) A revision of the genus *Inocybe* in Europe I. Subgenus *Inosperma* and the smooth-spored species of subgenus *Inocybe*. *Persoonia (Suppl.)* 3: 1-247.
- LUDWIG E (2017) *Pilzkompendium 4. Abbildungen. Cortinariaceae I – Cortinarius (part I), Galerina, Hebeloma, Hebelomina, Inocybe, Phaeogalera*. Fungicon Verlag. Berlin.
- MALENÇON G & BERTAULT R (1970) *Flore des champignons supérieurs du Maroc, Vol. 1*. Institut Scientifique Chérifien et Faculté des Sciences. Rabat. 604 pp.
- MATHENY PB, HOBBS AM & ESTEVE-RAVENTÓS F (2020) Genera of *Inocybaceae*: New skin for the old ceremony. *Mycologia* 112(1): 83-120. DOI: <https://doi.org/10.1080/00275514.2019.1668906>
- MOËNNE-LOCCOZ P, POIRIER J & REUMAUX P (1988) Compléments à la Deuxième partie des *Inocybes*. *Bull. Soc. Mycol. France* 104(4): Pl. 252-254, Addenda: 3-10.
- MUNSELL COLOR (1994) *Soil Color Charts (revised edition)*. Macbeth Division of Kollmorgen Instruments Corporation. New Windsor. New York. USA.
- REUMAUX P (1982) Miettes sur les *Inocybes*. *Doc. Mycol.* 12(48): 1-28.
- RONQUIST F, TESLENKO M, VAN DER MARK P, AYRES D, DARLING A, HOHNA S, LARGE B, LIU L & al. (2012) MrBayes 3.2: efficient Bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61: 539-554. DOI: <https://doi.org/10.1093/sysbio/sys029>
- STANGL J & VESELSKY J (1982) Risspilze der Sektion *Lilacinae* Heim (Beiträge zur Kenntnis seltener *Inocyben*. Nr. 19). *Ceska Mykol.* 36(2): 85-99.
- TRIFINOPOULOS J, NGUYEN L-T, VON HAESELER A & MINH BQ (2016) W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Res.* 44: W232-W235. DOI: <https://doi.org/10.1093/nar/gkw256>
- VELLINGA EC (1988) Glossary. In: BAS C, KUYPER TW, NOORDELOOS ME & VELLINGA EC (ed.). *Flora Agaricina Neerlandica vol. 1*: 54-64. AA Blakema. Rotterdam. The Netherlands.
- WHITE TJ, BRUNS T, LEE S & TAYLOR JW (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics: 315-322. In: INNIS MA, GELFAND DH, SNINSKY JJ & WHITE TJ (ed.). *PCR protocols: a guide to methods and applications*. Academic Press Inc., New York.